



DOI: 10.32768/abc.202293394-397



## Conservation Rates of Breast and Axilla Indicate the Quality of Surgical Management of the Early-Stage Breast Cancer Patient

Ana Car Peterko<sup>\*a</sup> , Damir Grebić<sup>a</sup> , Damir Juranić<sup>b</sup>, Aleksandra Bila<sup>a</sup>, Danijel Trčak<sup>a</sup> , Dolores Brajan<sup>a</sup>, Nevija Žepina<sup>a</sup>, Mladenka Golubovac Iskra<sup>a</sup>, Franjo Lovasić<sup>a</sup> 

<sup>a</sup>Department of General Surgery and Surgical Oncology, Clinical Hospital Centre Rijeka, Croatia

<sup>b</sup>Department of Plastic and Reconstructive Surgery, Clinical Hospital Centre Rijeka, Croatia

### ARTICLE INFO

**Received:**

21 February 2022

**Revised:**

13 March 2022

**Accepted:**

15 March 2022

**Keywords:**

Breast Centre, Certificate, quality indicator, mastectomy, breast-conserving

### ABSTRACT

**Background:** Multidisciplinary care in quality-assured specialized Breast Centre (BC) is nowadays considered optimal management of breast cancer and is associated with better patient outcomes. Modern breast cancer surgery should provide optimal oncological outcomes and preserve the quality of life. European Society of Breast Cancer Specialists (EUSOMA) set up the minimum requirements for a specialized BC and a set of 15 mandatory quality indicator (Q.I.) benchmarks for BC certification purposes. Six Q.I. are intended for breast cancer surgery quality assessment.

**Methods:** In BC of Clinical Hospital Centre Rijeka, data were collected retrospectively for 2019 and prospectively thereafter in a clinical register, encrypted according to the EUSOMA instructions and uploaded into collective EUSOMA database. Following database validation, all Q.I. were calculated for our BC for 3 consecutive years. In addition, a comprehensive on-site audit was performed in 2021 for all services included in breast cancer management in Rijeka.

**Results:** All mandatory surgical Q.I. were above the EUSOMA benchmarks in all 3 years. Non-compliance with EUSOMA recommendations were reported as major, minor, recommendations and observations. For BC surgical department, no major or minor non-conformities were reported. At the national level, mastectomy rates were above the permitted EUSOMA benchmark in 4 consecutive years.

**Conclusion:** We have voluntarily initiated and performed quality control of our BC. The certificate obtained is a confirmation of the high quality of care. Concerning the mastectomy rate in Croatia, other centres should consider quality evaluation to determine the status of detected suboptimal surgical management.

Copyright © 2022. This is an open-access article distributed under the terms of the [Creative Commons Attribution-Non-Commercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/) International License, which permits copy and redistribution of the material in any medium or format or adapt, remix, transform, and build upon the material for any purpose, except for commercial purposes.

### INTRODUCTION

Breast cancer is the most common malignant disease affecting females. One in ten women in Europe is diagnosed with breast cancer in her lifetime. Due to the implementation of national screening programs, as

well as improved women's health awareness, the majority of breast cancer cases are nowadays diagnosed in the early stage of the disease, thus leading to an excellent prognosis if managed appropriately. Multidisciplinary care in the quality-assured specialized Breast Centre (BC) is associated with the best oncological outcomes,<sup>1-3</sup> and is considered optimal management of breast cancer patients.<sup>4</sup> European Society of Breast Cancer Specialists (EUSOMA) set up the minimum requirements for a specialized BC,<sup>5</sup> as well as a set of 15 mandatory quality indicators (Q.I.)

**\*Address for correspondence:**

Ana Car Peterko, MD, breast surgeon and surgical oncologist  
Department of General Surgery and Surgical Oncology,  
Clinical Hospital Centre Rijeka, Croatia  
Tel: +385 51 658 111  
Email: [anacarpeterko@gmail.com](mailto:anacarpeterko@gmail.com)



for certification purposes and quality comparison between different centres.<sup>6</sup> BC's surgical department is defined with a minimum of 150 procedures for newly diagnosed breast cancers and at least 2 breast surgeons employed by the BC. In addition, 6/15 mandatory Q.I. evaluate surgical management, primarily the appropriateness of surgical indication and the quality of surgical technique. Every Q.I. represents a recommendation for the optimal management, and the associated benchmark is the permitted level of deviation from the clinical guidelines. This paper aims to report the quality of breast cancer surgical management in the first certified BC in Croatia, evaluating with six mandatory EUSOMA Q.I. in 3 consecutive years and comparing the data with the proposed benchmarks. The surgical Q.I. available at the national level was compared to the EUSOMA benchmark as well.

## METHODS

In BC of Clinical Hospital Centre (CHC) Rijeka, data were collected retrospectively for 2019 and prospectively thereafter in a clinical register, encrypted according to the EUSOMA instructions and uploaded into the collective EUSOMA database. Data collection in a prospective register was approved by the institutional ethics committee. Following database validation, all Q.I. were calculated for BC of CHC Rijeka for 3 consecutive years. In addition, a comprehensive on-site audit was performed in 2021 for all services included in breast cancer management in CHC Rijeka. Data of the Breast Working Group Registry, Croatian Society of Pathology (2017-2020) was used for the determination of the annual Q.I. 5 values at the national level.

## RESULTS

All mandatory surgical Q.I. calculated for BC of CHC Rijeka were above the EUSOMA benchmarks in all 3 consecutive years (Table 1).

Following the on-site audit, non-compliance with EUSOMA recommendations were reported and categorized as major, minor, recommendations and observations. For BC surgical department, three recommendations and one observation were recorded (Table 2). However, no major or minor non-conformities were noticed.

At the national level, calculated annual Q.I.5 values were significantly below the minimum of EUSOMA requirements in all 4 years (58%, 59%, 66% and 63%).

## DISCUSSION

Quality assessment represents the comparison of routine clinical practice with the evidence-based guidelines for optimal management. The quality of

surgery is determined by the appropriate selection of the optimal procedure, as well as by the quality of

**Table 1.** EUSOMA quality indicator in Clinical Hospital Center Rijeka

EUSOMA Quality Indicator (Q.I.)	Min (%)	target	2019	2020	2021
Q.I.5 Invasive Ca <=3cm treated with BCS	70	85	96	92	97
Q.I.6 In situ Ca <=2 cm treated with BCS	80	90	94	90	100
Q.I.7 DCIS with no axillary clearance	97	99	100	100	100
Q.I.10 Invasive Ca receiving just 1 operation	80	90	90	89	95
Q.I.11 DCIS receiving just 1 operation	70	90	78	100	91
Q.I.12 SLNB in cN0 invasive Ca	90	95	97	97	92

performance. Although several population-based studies suggest inferior survival for mastectomy patients when compared to breast conservation,<sup>7-12</sup> radical procedures, both in breast and axilla, are not considered inferior to the conservative approach in terms of oncological outcomes.<sup>13-15</sup> However, due to their significant morbidity,<sup>16</sup> and the adverse impact on the quality of life,<sup>17</sup> these procedures are not recommended for the management of early breast cancer patients. Modern breast cancer surgery should provide the highest level of quality of life for breast cancer survivors in addition to optimal oncological outcomes and mastectomy, with or without breast reconstruction, should no longer be offered as an option to the early-stage breast cancer patient without solid oncological indication. High rates of both mastectomy and axillary clearance (evaluated by Q.I. 5, 6, 7 and 12), indicate suboptimal surgical management, as well as the high rates of re-intervention following breast conservative surgery (evaluated by Q.I. 10 and 11), represent suboptimal surgical technique.

Herein, we have presented the results of quality evaluation of the surgical department at our BC in Rijeka, Croatia. The results are expressed as rates of EUSOMA proposed Q.I., calculated for our department in 3 consecutive years and compared with the defined benchmarks.

The results indicate continuous, high-quality surgical management of patients in BC of CHC Rijeka. Higher rates of breast conservation, compared to

**Table 2.** EUSOMA classification, recommendations, observation and positive aspects

Audit Report (8-9/7/2021)	Breast Centre Rijeka	Dpt. of Breast Surgery
Major non-conformities	7	0
Minor non-conformities	8	0
Recommendations	3/7	A medical photograph of the breast should be available to decide the best surgical strategy for every patient Breast Centers should collect data and yearly monitor the functional outcomes The Breast Centre should comply with Q.I. 14 (<=5 lymph nodes/SLNB) The audit team observed that the Breast Centre has not yet started to collect and analyze validated patient-reported outcomes (PROs)
Observations	1/6	using validated measurements (patient-reported outcome measures, PROMs). The audit team congratulates the surgeons on joint working arrangements and their desire to learn new techniques despite the pandemic
Positive aspects	1/3	

EUSOMA Q.I. 5 and 6 benchmarks, are achievable with oncoplastic techniques, with no adverse impact on oncological outcomes, patient satisfaction or re-intervention rates.<sup>18</sup>

However, Q.I.5, the only available Q.I. at the national level is continuously below the proposed benchmark, revealing suboptimal patient management in Croatia. The recognition of any deviation in clinical practice is the first step toward its improvement. Uniform comprehensive institutional and national databases are essential for quality control and identification of non-compliance with the guidelines. Additional clinical data are required to determine the status of sub-optimal breast cancer surgical management identified in Croatia.

### CONCLUSION

Implementation of quality assessment is one of the factors associated with better patient outcomes. We have voluntarily initiated and performed quality control of our BC. The certificate obtained is a confirmation of the high quality of care. Concerning the latest reports of breast cancer mortality rates in

Croatia,<sup>19</sup> as well as the high mastectomy rate in early-stage breast cancer surgery, all centres involved in breast cancer management should consider quality evaluation as well.

### ACKNOWLEDGEMENTS

We would like to thank Croatian Society of Pathology, regional coordinators of the Breast Working Group Registry: Snježana Tomić, Manuela Avirović, Melita Perić Balja, Marija Milković Periša, Valerija Blažičević, Čedna Tomasović, for data used to determine the annual Q.I. 5 values at the national level for the period from 2017 till 2020.

### CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

### ETHICAL CONSIDERATIONS

The protocol of the study was approved by the institutional ethics committee in a prospective register.

### FUNDING

Nothing to declare.

### REFERENCES

- Greenup RA, Obeng-Gyasi S, Thomas S, Houck K, Lane WO, Blitzblau RC, et al. The effect of hospital volume on breast cancer mortality. *Ann Surg* 2018;267(2):375e81. doi: 10.1097/SLA.0000000000002095.
- de Camargo Cancela M, Comber H, Sharp L. Hospital and surgeon caseload are associated with risk of re-operation following breast-conserving surgery. *Breast Canc Res Treat* 2013;140(3):535e44. doi: 10.1007/s10549-013-2652-5.
- Gooiker GA, van Gijn W, Post PN, van de Velde CJ, Tollenaar RA, Wouters MW. A systematic review and meta-analysis of the volume outcome relationship in the surgical treatment of breast cancer. Are breast cancer patients better off with a high volume provider? *Eur J Surg Oncol* 2010;36(Suppl 1):S27e35. doi: 10.1016/j.ejso.2010.06.024.
- Cataliotti L, Costa A, Daly PA, Fallowfield L, Freilich G, Holmberg L, et al. Florence statement on breast cancer: forging the way ahead for more research on and better care in breast cancer. *Eur J Canc* 1999;35(1):14-5. doi: 10.1016/S0959-8049(98)00384-0.
- Biganzoli L, Cardoso F, Beishon M, Cameron D, Cataliotti L, Coles CE, et al. The requirements of a



- specialist breast centre. *Breast*. 2020 Jun;51:65-84. doi: 10.1016/j.breast.2020.02.003.
6. Biganzoli L, Marotti L, Hart CD, Cataliotti L, Cutuli B, Kühn T, et al. Quality indicators in breast cancer care: An update from the EUSOMA working group. *Eur J Cancer*. 2017 Nov;86:59-81. doi: 10.1016/j.ejca.2017.08.017.
  7. Hartmann-Johnsen OJ, Kåresen R, Schlichting E, Nygård JF. Better survival after breast-conserving therapy compared to mastectomy when axillary node status is positive in early-stage breast cancer: a registry-based follow-up study of 6387 Norwegian women participating in screening, primarily operated between 1998 and 2009. *World J Surg Oncol*. 2017 Jul 3;15(1):118. doi: 10.1186/s12957-017-1184-6.
  8. Hartmann-Johnsen OJ, Kåresen R, Schlichting E, Nygård JF. Survival is Better After Breast Conserving Therapy than Mastectomy for Early Stage Breast Cancer: A Registry-Based Follow-up Study of Norwegian Women Primary Operated Between 1998 and 2008. *Ann Surg Oncol*. 2015 Nov;22(12):3836-45. doi: 10.1245/s10434-015-4441-3.
  9. Nandakumar A, Rath GK, Kataki AC, Bapsy PP, Gupta PC, Gangadharan P, et al. Decreased Survival With Mastectomy Vis-à-Vis Breast-Conserving Surgery in Stage II and III Breast Cancers: A Comparative Treatment Effectiveness Study. *J Glob Oncol*. 2016 Oct 12;3(4):304-313. doi: 10.1200/JGO.2016.004614.
  10. Agarwal S, Pappas L, Neumayer L, Kokeny K, Agarwal J. Effect of breast conservation therapy vs mastectomy on disease-specific survival for early-stage breast cancer. *JAMA Surg*. 2014 Mar;149(3):267-74. doi: 10.1001/jamasurg.2013.3049.
  11. Christiansen P, Carstensen SL, Ejlersen B, Kroman N, Offersen B, Bodilsen A, et al. Breast conserving surgery versus mastectomy: overall and relative survival-a population based study by the Danish Breast Cancer Cooperative Group (DBCG). *Acta Oncol*. 2018 Jan;57(1):19-25. doi: 10.1080/0284186X.2017.1403042.
  12. Lagendijk M, van Maaren MC, Saadatmand S, Strobbe L, Poortmans P, Koppert LB, et al., "Breast conserving therapy and mastectomy revisited: breast cancer specific survival and the influence of prognostic factors in 129,692 patients," *International Journal of Cancer* 2018, vol. 142, no. 1, pp. 165–175. doi: 10.1002/ijc.31034.
  13. Veronesi U, Cascinelli N, Mariani L, Greco M, Saccozzi R, Luini A, et al. Twenty-year follow-up of a randomized study comparing breast-conserving surgery with radical mastectomy for early breast cancer. *N Engl J Med*. 2002 Oct 17;347(16):1227-32. doi: 10.1056/NEJMoa020989.
  14. Veronesi U, Luini A, Galimberti V, Zurrada S. Conservation approaches for the management of stage I/II carcinoma of the breast: Milan Cancer Institute trials. *World J Surg*. 1994 Jan-Feb;18(1):70-5. doi: 10.1007/BF00348194.
  15. Fisher B, Anderson S, Bryant J, Margolese RG, Deutsch M, Fisher ER, et al. Twenty-year follow-up of a randomized trial comparing total mastectomy, lumpectomy, and lumpectomy plus irradiation for the treatment of invasive breast cancer. *N Engl J Med*. 2002 Oct 17;347(16):1233-41. doi: 10.1056/NEJMoa022152.
  16. Chand ND, Browne V, Paramanathan N, Peiris LJ, Laws A, Rainsbury RM. Patient-Reported Outcomes Are Better after Oncoplastic Breast Conservation than after Mastectomy and Autologous Reconstruction. *Plast Reconstr Surg Glob Open*. 2017 Jul 24;5(7):e1419. doi: 10.1097/GOX.0000000000001419.
  17. Potter S, Trickey A, Rattay T, O'Connell RL, Dave R, Baker E, et al. Therapeutic mammoplasty is a safe and effective alternative to mastectomy with or without immediate breast reconstruction. *BJS* 2020; 107(7):832-844. doi: 10.1002/bjs.11468.
  18. Losken A, Dugal CS, Styblo TM, Carlson GW. A meta-analysis comparing breast conservation therapy alone to the oncoplastic technique. *Ann Plast Surg*. 2014 Feb;72(2):145-9. doi: 10.1097/SAP.0b013e3182605598.
  19. Dafni U, Tsourti Z, Alatsathianos I. Breast Cancer Statistics in the European Union: Incidence and Survival across European Countries. *Breast Care (Basel)*. 2019 Dec;14(6):344-353 doi: 10.1159/000503219.

### How to Cite This Article

Peterko AC, Grebić D, Juranić D, Bila A, Trčak D, Brajan D, et al. Conservation Rates of Breast and Axilla Indicate the Quality of Surgical Management of the Early-Stage Breast Cancer Patient. *Arch Breast Cancer*. 2022; 9(3): 394-7.

Available from: <https://www.archbreastcancer.com/index.php/abc/article/view/553>